


Prepared in cooperation with the Bureau of Land Management



## Mapping Evapotranspiration Units in the Basin and Range Carbonate-Rock Aquifer System, White Pine County, Nevada, and Adjacent Areas in Nevada and Utah



Scientific Investigations Report 2007–5087

# **Mapping Evapotranspiration Units in the Basin and Range Carbonate-Rock Aquifer System, White Pine County, Nevada, and Adjacent Areas in Nevada and Utah**

By J. LaRue Smith, Randell J. Laczniak, Michael T. Moreo, and Toby L. Welborn

Prepared in cooperation with the Bureau of Land Management

Scientific Investigations Report 2007-5087

**U.S. Department of the Interior**  
**U.S. Geological Survey**

**U.S. Department of the Interior**  
DIRK KEMPTHORNE, Secretary

**U.S. Geological Survey**  
Mark D. Myers, Director

U.S. Geological Survey, Reston, Virginia: 2007

For product and ordering information:

World Wide Web: <http://www.usgs.gov/pubprod>

Telephone: 1-888-ASK-USGS

For more information on the USGS--the Federal source for science about the Earth, its natural and living resources, natural hazards, and the environment:

World Wide Web: <http://www.usgs.gov>

Telephone: 1-888-ASK-USGS

Any use of trade, product, or firm names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

Although this report is in the public domain, permission must be secured from the individual copyright owners to reproduce any copyrighted materials contained within this report.

Suggested citation:

Smith, J.L., Laczniak, R.J., Moreo, M.T., and Welborn, T.L., 2007, Mapping evapotranspiration units in the Basin and Range carbonate-rock aquifer system, White Pine County, Nevada, and adjacent areas in Nevada and Utah: U.S. Geological Survey Scientific Investigations Report 2007-5087, 20 p.

Nichols (2001) mapped areas of ground-water discharge based on the outer boundary of phreatophytes. Nichols mapped these boundaries by using basic reconnaissance-type field methods. His mapping generally was done in early to mid-summer when contrast in the color of greasewood and other rangeland vegetation is high. Control points along roads or trails typically were established by using Global Positioning System (GPS) equipment. Satellite data, aerial photography, and elevation models were used along with these established control points to map the complete phreatophyte boundary at a scale of 1:24,000. This general mapping technique also was used by Laczniaik and others (2001) in developing ground-water discharge estimates for the Death Valley regional ground-water flow system. Digital maps delineating the outer boundary of phreatophyte areas throughout central Nevada developed by Nichols (2001) are available in Smith and others (1999) and were acquired for this study.

The Southern Nevada Water Authority (SNWA) mapped areas of ground-water discharge in selected hydrographic areas of eastern Nevada in 2004 (Michael Wallen, Southern Nevada Water Authority, oral commun., 2006). Boundaries previously mapped by Nichols (2001) and aerial photography and other available GIS data were used to guide field and mapping efforts. These boundaries were acquired by the USGS for use in this study.

The boundaries established by Harrill and others (1988), Nichols (2001), and the SNWA all were used to guide field activities and mapping efforts directed at delineating potential areas of ground-water discharge in the study area. The boundaries mapped by these previous investigations were integrated and compared. Identified discrepancies were resolved by field verification efforts conducted in mid-July 2005. The final boundary identifies the outer extent where shallow ground water may be consumed by ET. The area within this boundary is referred to as a potential area of ground-water discharge and the area outside the boundary is assumed to be an area of no ground-water discharge. The potential area of ground-water discharge within a hydrographic area can be a single area delineated by one continuous boundary or can be multiple areas. The size and number of these areas within a hydrographic area is independent of valley size.

Many of the valleys in the study area are large and often difficult to visit because accessible roads and trails are lacking. A helicopter survey was used to map phreatophyte boundaries in selected valleys during a 2-day period in June 2005. The helicopter was equipped with GPS equipment to record flight-path coordinates. The helicopter flew at an altitude of about 100 ft above land surface and at a speed of about 80 mi/h. Mapping of phreatophytes such as greasewood at this time of year from the air is straightforward, considering the greenness

of greasewood compared to the many nonphreatophytic species present in the area. A video clip ([clip 1](#)) of a flight through Snake Valley illustrates the contrast and visibility of greasewood in a discharge area typical of the study area. The video clip begins with the helicopter flying northward with the greasewood shrubs on the right and the rangeland grasses on the left. Near the end of the clip the helicopter follows the greasewood boundary by making a left turn. The potential areas of ground-water discharge used to calculate ET in the study were developed by combining and integrating phreatophyte boundaries delineated in previous studies, numerous field mapping efforts, and the phreatophyte extents mapped by the helicopter survey. Potential areas of ground-water discharge as mapped for the study are shown in [figure 3](#). Specific data sources used to delineate the potential areas of ground-water discharge within each hydrographic area are reported in [table 2](#).


## Delineation of Evapotranspiration Units

An ET-unit map was created by using data from SWReGAP ecological systems, interpreted Landsat data from multiple dates, and Landsat MSAVI and Tasseled Cap products. An explanation of the data used, the processing techniques, and a field accuracy assessment are described in the following section.

## Southwest Regional Gap Analysis Program

The SWReGAP identifies 125 land-cover classes referred to as ecological systems. An ecological system is a mid-scale classification unit defined by NatureServe <http://www.natureserve.org/publications/usEcologicalsystems.jsp>. Ecological systems represent recurring groups of biological communities that exist in similar physical environments. Selected classes of the SWReGAP that conform with characteristics of defined ET units were used in developing ET-unit distributions. The ET units defined by SWReGAP classes are marshlands, dry playa, and open water bodies. The mapping of these SWReGAP classes relied on multiple date imagery over a season and is consistent with the NLCD. Other ecological systems also were evaluated as to whether they adequately represented the extent of other ET units. Other evaluated systems included mixed salt desert scrub, greasewood flat, and big sagebrush shrubland. Field observations indicated that these ecological systems did not correlate well with ET units and tended to map an area larger than the outer extent of potential areas of ground-water discharge.

8 Mapping Evapotranspiration Units, Basin and Range Carbonate-Rock Aquifer System, Nevada and Utah



Base from U.S. Geological Survey digital data 1:100,000, 1978-89. Universal Transverse Mercator Projection, Zone 11, NAD 83. Shaded relief base from 1:250,000-scale Digital Elevation Model; sun illumination from northwest at 30 degrees above horizon.

**EXPLANATION**

- Potential areas of ground-water discharge
- Boundary of study area
- Boundary of hydrographic area
- Boundary of subbasin and No.

**Figure 3.** Potential areas of ground-water discharge in the Basin and Range carbonate-rock aquifer system study area, Nevada and Utah.

**Table 2.** Potential areas of ground-water discharge mapped by hydrographic area in the Basin and Range carbonate-aquifer system study area, Nevada and Utah.

[**Helicopter:** Aerial mapping done during 2005 study using helicopter.  
**Southern Nevada Water Authority (SNWA):** Field verified column indicates that hydrographic area was visited and that the boundary was checked by field personnel in 2005 (Michael Wallen, unpub. data, 2005)]

Hydrographic area	Method or Report				Field verified
	Helicopter	SNWA	Nichols (2001)	Harrill and others (2005)	
Butte Valley		×	×	×	×
Cave Valley	×	×			
Jakes Valley		×	×		×
Lake Valley	×			×	×
Little Smoky Valley			×	×	×
Long Valley		×	×	×	×
Newark Valley			×	×	×
Snake Valley	×	×		×	
Spring Valley		×	×	×	×
Steptoe Valley		×	×	×	
Tippett Valley			×		
White River Valley	×	×		×	×

## Landsat Thematic Mapper Imagery

Landsat Thematic Mapper imagery was used to delineate ET units in the study area. Landsat satellites orbit the Earth such that they acquire data over the same area at the same time every 16 days. These Earth-orbiting satellites are equipped with sensors to detect and acquire solar-reflected and earth-emitted radiation. Spectral reflectance, as acquired by Landsat sensors, represents an average value over a pixel (picture element) that measures about 100 ft by 100 ft. These pixel dimensions define the spatial resolution of the imagery.

Landsat data are available as scenes that image an area measuring about 115 mi by 115 mi. Each scene is defined by a path and row number. Multi-scene units are defined as consecutive scenes (rows) within a single path acquired and archived on the same date. Two multi-scene units were acquired to delineate ET units in the study area: (1) entity-id It5039032000019419 inclusive of scenes acquired July 12, 2005, along path 39 at rows 32, 33 and 34; and (2) entity-id It5040032000018400 inclusive of scenes acquired July 3, 2005, along path 40 at rows 32, 33, and 34. These scenes were selected because they met the criteria required to best delineate ET units. Scenes were contemporary with other ongoing field and data-collection activities being conducted as part of the

study. Experience based on recently completed ET studies indicates that the optimum acquisition dates for delineating ET units in the study area fall between late June and early July—June 21 being the day with the longest daylight hours (a period of peak solar radiation), and early July, the approximate period when transpiration by phreatophytes is near its peak. Another factor considered in image selection is the potential influence of any recent precipitation. Snowpack telemetry (SNOTEL) sites operated by the Natural Resources Conservation Service were reviewed at Ward Mountain, located south of Ely and Berry Creek, just east of Ely (fig. 2). On the basis of data collected from these telemetry sites, no significant precipitation had occurred since June 21, and therefore, vegetative conditions captured in these images are assumed not to be influenced by local precipitation.

All acquired imagery was georeferenced to allow geospatial evaluations and direct comparison with other spatially referenced datasets. Georeferencing assigns map coordinates to an image by using known ground control points and the corresponding image row and column. A mathematical transformation equation is determined to relate the image to map coordinates (Lillesand and Kiefer, 1987). Previously georeferenced Landsat imagery by the USGS that was acquired in 2000 was used in the georeferencing process. Corrections to georeference the 2005 images were applied by using a first-order polynomial equation. Each 2005 multi-scene image used between 30 and 35 control points. The root mean square error or positional accuracy for each multi-path image was about 100 ft.

Image standardization is the process of normalizing imaged spectral data for differences in sun illumination geometry, atmospheric effects, and instrument calibration. Standardization is required to accurately compare pixel spectral reflectance between images of different dates. Pixel values in each image were standardized by using software developed and distributed by the Remote Sensing and GIS Laboratory of Utah State University <http://www.gis.usu.edu/docs/projects/swgap/ImageStandardization.htm>. For atmospheric corrections, the technique COST without Tau was used. The interested reader is directed to Huang and others (2007) for more detail on the normalization procedure.

## Recently Irrigated Cropland

Recently irrigated cropland was delineated for the study area by Welborn and Moreo (2007). These investigators digitized cropland areas by using Landsat imagery from 2000, 2002, and 2005. Digitized field delineations were stored as polygons in an ArcGIS geodatabase and attributed with information including water source, method of application, and crop type. Each digitized field was field checked and verified during the summer of 2005.

### Modified Soil Adjusted Vegetation Index and Tasseled Cap Transformation

ET units not mapped by SWReGAP were delineated by using the Modified Soil Adjusted Vegetation Index (MSAVI) (Qi and others, 1994) and a Tasseled Cap transformation (Huang and others, 2002) from normalized Landsat data. A vegetation index is a number that is generated by the combination of remote sensing bands and has some relation to the amount of vegetation in a given pixel. The MSAVI is an index which removes soil influences from the vegetation index at sparse plant cover. Because sparsely vegetated conditions are most important in the Great Basin, the MSAVI is considered an appropriate index from which to map changes in plant cover. The Tasseled Cap transformation is being used to develop land-cover classes for the NLCD. The Tasseled Cap transformation compresses all the satellite spectral bands into a few bands associated with physical scene characteristics of brightness, greenness, and wetness. Brightness is a weighted sum of all Landsat reflectance bands, which is a response to changes in total reflectance, driven primarily by soil reflectance changes. Greenness contrasts the sum of the visible and near-infrared bands and has been shown to correlated to percentage canopy coverage. Wetness contrasts the sum of the visible and near-infrared bands against the sum of the longer wavelength bands (Crist and Cicone, 1984).

The MSAVI was used to map phreatophytic shrubs and grasses in the potential areas of ground-water discharge. Vegetation indices have been used often to quantify the abundance and vigor of vegetation imaged by multispectral sensors (U.S. Geological Survey EROS Data Center, 2006). The use of vegetation indices to map vegetation cover is based on the assumption that the greener and denser the vegetation, the greater the vegetation index. Nichols (2001) takes advantage of this relation and uses the MSAVI to quantify the amount of ground water lost to the atmosphere by ET. In applying this approach, Nichols assumes that ground-water ET is a function of ET, and that ET is a function of plant cover. Nichols (2001, p. B8) uses MSAVI to develop ranges in plant cover and describes the type or types of vegetation occurring in each defined plant-cover range. In general, shrubs dominate where plant cover is less than 35 percent and grasses dominate where plant cover is greater than 35 percent. Xerophyte conditions, areas where phreatophytes are absent and bare soil dominates, exist where plant cover is less than about 7 percent. These general relations between percentage of cover and plant type also were observed in the study area during field visits. ET units were defined by using breaks in MSAVI values to correlate with boundaries between xerophyte areas (no substantial ground-water discharge) and phreatophyte areas dominated by sparse desert shrubland, moderately dense desert shrubland, dense desert shrubland, grassland, or meadowland.


MSAVI values are dimensionless and range from -1.0 to 1.0. For this study, values from 0 to 1 were scaled from 0 to 200. Values less than 0 were set to 0. Values 0 to 14 were identified as xerophyte, 15 to 20 are sparse desert shrubland, 21 to 28 are moderately dense desert shrubland, 29 to 43 are dense desert shrubland, 44 to 55 are grassland, and 56 to 92 are meadowland. The MSAVI values from 93 to 200 usually occurred where ET units were assigned to recently irrigated cropland or marshland. If not, they were assigned to meadowland.

The Tasseled Cap brightness, greenness, and wetness bands were used to map areas of moist bare soil. An unsupervised classification based on these three bands was applied to the data set. Unsupervised classification clusters pixels in a data set according to statistics only, without any user-defined training classes. The user then identifies clusters that relate to what is being mapped. Yelland Dry Lake in northern Spring Valley and an area around Twin Springs in northern Snake Valley (fig. 2) were used to identify the moist bare soil. These areas of moist bare soil are intended to represent locations where moisture is present throughout much of the year and where the depth to ground water is shallow.

The Tasseled Cap band of greenness also was considered in mapping shrubs and grasses. The SNWA established shrub transects in different hydrographic areas of the study area. These shrub transects defined the occurrence of shrubs over a 328-foot linear length of vegetation. Forty-five of these were used to define the percentage of plant cover. Values of the Tasseled Cap greenness and MSAVI at each transect were compared to the percentage of plant cover. The MSAVI showed a slightly higher correlation than the Tasseled Cap (fig. 4). The correlation of the MSAVI and the Tasseled Cap greenness with the percent plant cover is similar, but the MSAVI correlates better with the lower percentage of plant cover. The MSAVI was chosen to map the ET units representing shrubs and grasses.

### Assembly and Distribution of Evapotranspiration Units

A map presenting the spatial distribution of ET units was developed by using units defined by the MSAVI and Tasseled Cap analysis, the delineation of recently irrigated cropland, and the data of SWReGAP products. A single ET-unit map was created from ET units developed by each of these different data sets by layering the data according to priorities. The initial ET-unit map was assembled by first using the MSAVI data set. The moist bare soil ET unit, derived by the Tasseled Cap procedure, replaced the MSAVI data wherever moist bare soil occurred. The ET units of marshland, open water, and dry playa, derived from the SWReGAP data, replaced the previous



**Figure 4.** Percentage of plant cover determined from transects compared to scaled values of (A) Tasseled Cap greenness band and (B) modified soil adjusted vegetation index.

MSAVI and Tasseled Cap data wherever those ET units occurred. Finally, the recently irrigated cropland, derived from photo interpretation of the three Landsat data sets, replaced ET units from the other data sets wherever recently irrigated cropland occurred. The ET units and the source of their information are summarized in [table 3](#).

**Table 3.** Evapotranspiration (ET) units and the source used to develop the ET-unit map.

ET unit	Source
Open water, marshland, dry playa	SWReGAP
Recently irrigated cropland	Landsat 2000, 2002, 2005
All shrubland, grassland, and meadowland	MSAVI from 2005
Moist bare soil	Tasseled Cap from 2005
Xerophyte	MSAVI from 2005

A filtering technique was applied to all ET units except those representing recently irrigated cropland and those developed from SWReGAP. The filtering technique added and removed pixels on the basis of a minimum mapping unit. The size of the minimum mapping unit depends on the scale and resolution of the spectral data set being used to delineate land cover (Anderson and others, 1976) and on the intended use of the map product. The processing of SWReGAP by the Gap Analysis Program included a minimum mapping unit filter that was set to 1.1 acres as described in the SWReGAP for Nevada, Utah, New Mexico, Colorado, and Arizona Project Status (U.S. Environmental Protection Agency, 2007). The procedure is described in section, "Mapping Methods and Accuracy" of the Southeast Idaho Mapping Zone Documentation (U.S. Geological Survey, 2007a). This same procedure and criteria were used to filter the ET units defined by the MSAVI and Tasseled Cap. The minimum mapping unit was set at four adjacent pixels which measure about 35,000 ft<sup>2</sup> or 0.8 acre in the source imagery. An area of this size provides sufficient resolution within the fetch areas of ET stations established as part of the study (Moreo and others, 2007). The ET units as mapped within the delineated potential areas of ground-water discharge by using the filtering technique and the methods and priorities previously discussed are shown in [figure 5](#).

The shrubland units are the most prominent ET unit in the study area ([fig. 6](#)). The three shrubland ET units account for about 83 percent of the acreage delineated as potentially contributing to ground-water discharge. The more-densely vegetated meadowland and marshland ET units typically occur near springs and along major spring-drainage channels near the center of the valley floor, whereas the less-densely vegetated ET units typically occur along the outer edges of discharge areas ([fig. 5](#)). The more-densely vegetated ET units of marshland, meadow, and grassland account for only about 6 percent of the total delineated acreage, whereas, the least-vegetated units (dry playa, moist bare soil, sparse desert shrubland, moderately dense desert shrubland, and dense desert shrubland) account for about 91 percent. About 1,700 mi<sup>2</sup> are covered with phreatophytes or moist bare soil in the study area and potentially discharge ground water. The delineated acreage of each ET unit is shown by major hydrographic area in [table 4](#).

ET-unit acreage also is provided in [table 4](#) by subbasin. Subbasins are internal divisions within a hydrographic area. Major hydrographic areas in the study area were subdivided into these smaller internal units to help spatially resolve and refine the current understanding of interbasin flow. Subbasins are delineated such that their boundaries represent potential hydrologic divides within the larger hydrographic area and are based primarily on geophysical interpretations of gravity data


and assumed properties of known subsurface rock (Sweetkind and others, 2007). Subbasin boundaries commonly were delineated to coincide with interpreted intrabasin shallow bedrock. As presented in [table 4](#), the subbasins within a hydrographic area are not named but identified numerically ([fig. 2](#)). The first subbasin in a hydrographic area is the northernmost area. If a hydrographic area is not divided into subbasins, it is identified as subbasin 1.

## Evapotranspiration-Unit Field Accuracy

An assessment of the mapped accuracy of the ET units provides information to potential users about the maps reliability and usefulness as related to some intended purpose. Some needs require high-accuracy maps, whereas others may require much less spatial accuracy. Map quality and accuracy typically are assessed by using a probability based sampling design and relatively large number of samples per class or map unit. This type of assessment often is referred to as a map accuracy assessment. The quality of a map's assessed accuracy depends on the quality of the information used to ground truth or verify the map data. For the intended purpose of this map product, field-based ground-truth observations were considered sufficient to assess accuracy. A discussion of map accuracy, methods, and protocol can be found at U.S. Geological Survey (2007b).

Many different sources can introduce bias into an accuracy assessment. For example, bias can be introduced by sampling only along roads that provide easy access. The accuracy of a mapped ET unit can be affected by the nature of its boundary. For example, transitional boundaries are more difficult to locate, whereas boundaries defined by an abrupt change from one unit to the next are more easily detected. Because the ET-unit map has sharp and distinguished boundaries to delineate between different ET units, whereas in nature, different ET units transition spatially, an ET-unit map does not reflect this transitional nature. The ET-unit map rather provides estimates of acreages associated with generalized vegetation and/or soil-moisture conditions from which to make estimates of evapotranspiration.

Field assessment of the accuracy of the shrubland, grassland, meadowland and xerophyte ET units in Snake and Spring Valleys was done during August and September of 2006. In Snake Valley 28 field sites were visited and in Spring Valley 38 were visited. The guidelines used to direct field activities closely followed those described by the SWReGAP in the "Field Methodologies and Training Manual for Nevada Field Crews" document (U.S. Environmental Protection Agency, 2003).




Base from U.S. Geological Survey digital data 1:100,000, 1978-89. Universal Transverse Mercator Projection, Zone 11, NAD 83. Shaded relief base from 1:250,000-scale Digital Elevation Model; sun illumination from northwest at 30 degrees above horizon.

**EXPLANATION**

- |   |                    |   |                                   |
|---|--------------------|---|-----------------------------------|
|  | Open water         |  | Dense desert shrubland            |
|  | Irrigated cropland |  | Moderately dense desert shrubland |
|  | Marshland          |  | Sparse desert shrubland           |
|  | Meadowland         |  | Moist bare soil                   |
|  | Grassland          |  | Dry playa                         |
- 
-  Boundary of study area
  -  Boundary of hydrographic area
  -  Boundary of subbasin and No.

**Figure 5.** Spatial distribution of evapotranspiration (ET) units in the Basin and Range carbonatite-rock aquifer system study area, Nevada and Utah.



**Figure 6.** Evapotranspiration (ET)-unit acreage and percentage in the Basin and Range carbonate-rock aquifer system study area, Nevada and Utah.

The characterization of shrub-dominated environments requires an estimate of the approximate area of vegetation cover. Making this estimate whether by observation or by measurement can be difficult where plant vigor, dimensions, and spacing vary. To help guide this effort, field personnel viewed charts which have been used to estimate the percentage composition of rocks and sediments (Compton, 1962, p. 332-333). With these charts, the black specs were considered to be shrubs in order to estimate the percentage of plant cover within the circle. Estimates of plant cover based on these charts are expected to be accurate to within about 5 percent.

A single evaluation of how well the field crew estimated vegetation cover by using the geology field charts was done near the Snake Valley ET Station. An estimate of the percentage of vegetation cover was made by performing an unsupervised classification (as explained in the “Modified Soil Adjusted Vegetation Index and Tasseled Cap Transformation” section of this report) from a photograph taken with a hand-held digital camera during the helicopter survey. The photographic image covered an area of about 50 by 40 ft with a spatial resolution of approximately 0.3 in. and is shown in [figure 7A](#). An unsupervised classification of this photograph produced 32 different clusters. These clusters were combined into green plant leaf, leafless wood, soil, and shadow groups. This was done by identifying those clusters that were green colors on the photograph which represented green vegetation, for example. This also was done for the various soil colors, dark shadows, and leafless wood. The resulting percentage


of each group was 13.1 for green plant leaf, 4.8 for leafless wood, 60.2 for soil, and 21.9 for shadows ([fig. 7B](#)). Some shadowed area is likely to include active greasewood. Non-green leafed wood is not considered shrub cover in that it does not contribute to ET and is not characterized as vegetation by vegetation indexes computed from remotely sensed spectral data. Assuming that about one tenth of the shadowed area is active vegetation, shrub cover for this area is estimated at about 15 percent. This estimate was consistent with the 10- to 20-percent moderately dense desert shrubland estimated by field observation at this site using the geology field charts to estimate percentage of vegetation cover (Moreo and others, 2007, table 3, p. 13).

Map accuracy can be expressed as overall accuracy, producer accuracy, and user accuracy. To determine map accuracies for this study, the ET unit identified from the field observations was compared to the ET unit of the map. Overall accuracy is the most general measure and is calculated as the number of correct observations divided by the total number of observations. This does not mean that every ET unit was successfully classified at that accuracy. The producer’s accuracy relates to the probability that a field visited ET unit will be correctly mapped on the ET-unit map and measures the errors of omission. In contrast, the user’s accuracy indicates the probability that a sample from the ET-unit map actually matches what is found at the field visited sites and measures the error of commission (U.S. Geological Survey, 2007b).

**Table 4.** Evapotranspiration (ET)-unit acreage delineated by hydrographic area and subbasin in Basin and Range carbonate-rock aquifer system study area, Nevada and Utah.

[Hydrographic area subbasin: SB is subbasin. Number is subbasin number. Active ET-unit total does not include xerophytic acreage. –, not applicable]

Hydro-graphic-area sub-basin	ET-unit acreage, in acres											Active ET-unit total
	Xero-phytic	Marsh-land	Meadow-land	Grass-land	Dense desert shrubland	Moderately dense desert shrubland	Sparse desert shrubland	Moist bare soil	Open water	Dry playa	Irrigated cropland	
Butte Valley												
1	141	64	478	615	5,827	50,897	7,891	0	4	0	202	65,977
2	8	0	0	0	79	3,244	371	0	0	0	0	3,694
Total	149	64	478	615	5,906	54,141	8,262	0	4	0	202	69,671
Cave Valley												
1	2	81	503	280	842	354	6	0	0	0	0	2,066
2	141	0	0	2	534	7,005	3,546	0	0	194	0	11,282
Total	143	81	503	283	1,376	7,359	3,552	0	0	194	0	13,348
Jakes Valley												
–	27	25	91	146	540	203	6	0	26	0	187	1,224
Lake Valley												
1	962	630	1,143	822	4,077	32,384	16,296	0	26	94	0	55,472
2	0	0	0	0	0	0	0	0	0	0	0	0
Total	962	630	1,143	822	4,077	32,384	16,296	0	26	94	0	55,472
Little Smoky Valley												
Northern part	409	62	355	379	1,191	1,678	2,108	0	0	5	216	5,994
Central part	0	0	0	0	0	0	0	0	0	0	0	0
Total	409	62	355	379	1,191	1,678	2,108	0	0	5	216	5,994
Long Valley												
–	78	2	3	4	1,219	12,155	4,901	0	0	0	0	18,283
Newark Valley												
1	5,218	996	2,247	1,397	6,228	11,110	2,556	1	1	1,111	208	25,856
2	8,606	192	639	661	3,620	14,284	6,035	0	1	10,625	285	36,341
3	110	0	1	0	172	7,526	2,830	0	0	24	0	10,553
Total	13,933	1,188	2,887	2,058	10,020	32,920	11,421	1	2	11,760	493	72,750
Snake Valley												
1	8,603	334	693	347	2,527	6,854	17,772	0	0	56,499	1,785	86,810
2	16,034	541	1,746	1,463	7,988	34,568	66,023	578	115	5,553	1,138	119,711
3	3,896	432	1,696	799	3,638	26,758	46,090	0	100	1,081	5,136	85,729
4	92	535	1,816	834	7,368	17,297	3,254	0	212	0	1,873	33,189
5	0	0	0	0	0	0	0	0	0	0	0	0
Total	28,625	1,842	5,951	3,443	21,520	85,476	133,139	578	427	63,133	9,932	325,440
Spring Valley												
1	1	119	303	154	747	377	61	0	0	0	0	1,761
2	5,761	1,259	3,223	2,386	13,055	43,870	23,563	2,810	6	15,509	2,867	108,549
3	1,239	699	1,639	1,007	9,301	39,639	12,083	9	0	520	2,492	67,389
4	0	0	0	0	0	0	0	0	0	0	0	0
Total	7,001	2,076	5,165	3,547	23,104	83,886	35,707	2,819	6	16,029	5,359	177,698
Steptoe Valley												
1	8,762	790	2,251	2,442	16,691	69,506	26,861	237	5	2,464	2,766	124,013
2	968	2,993	5,171	3,813	13,197	15,992	3,158	21	147	0	2,354	46,848
3	17	152	498	254	1,427	985	76	0	287	0	0	3,679
Total	9,746	3,936	7,920	6,509	31,315	86,483	30,096	258	440	2,464	5,120	174,540
Tippet Valley												
–	42	0	21	51	1,013	4,569	1,623	0	0	497	0	7,775
White River Valley												
1	46	142	340	188	737	748	134	0	0	0	841	3,131
2	239	48	293	253	2,419	15,552	9,881	0	0	19	490	28,954
3	27	104	1,114	1,083	4,733	4,953	541	0	0	9	4,965	17,503
4	2,006	2,877	2,182	2,413	16,450	68,298	34,353	14	685	941	295	128,508
Total	2,318	3,172	3,929	3,936	24,340	89,551	44,909	14	685	970	6,591	178,096
Basin and Range carbonate-aquifer system study area												
Total		13,079	28,444	21,793	125,620	490,805	292,020	3,670	1,616	95,145	28,101	1,100,292



**EXPLANATION**

**Image B color-based land cover class**

	Green plant leaf (13.1 percent)		Leafless wood (4.8 percent)
	Shadow (21.9 percent)		Soil (60.2 percent)

**Figure 7.** (A) natural color and (B) color-classified photograph of greasewood near ET station in Snake Valley, Nevada. Photograph taken looking downward from helicopter during aerial survey, June 6, 2005.

Accuracies for the SWReGAP ecological systems were computed qualitatively. A detailed description of these accuracies can be obtained from the SWReGAP 'PROVISIONAL' Landcover and Related Datasets document (Southwest Regional Gap Analysis Project, 2007a). Accuracies are presented as summaries that are brief descriptions provided by the teams involved in the mapping process. Summaries are intended to provide an evaluation on how well each class was mapped from the perspective of the land-cover mapping analyst, taking into consideration the number of training and reference samples in the observation sample set for the cover class and the team's knowledge and

familiarity with the mapping area. Two SWReGAP mapping zones, NV-4 and UT-1, are present in the study area. The qualitative summaries describing the accuracy of ecological systems representing open water, marshland, and dry playa ET units mapped in zones NV-4 and UT-1 are briefly summarized in [table 5](#). The overall accuracy of the SWReGAP for all ecosystems in NV-4 and UT-1 is reported to be 53 and 59 percent, respectively (Southwest Regional Gap Analysis Project, 2007b). [Table 5](#) also provides qualitative summaries for the desert shrubland, grassland, and meadowland ET units mapped as part of the study and based on field work described in the previous paragraph.

**Table 5.** Field accuracy assessment of evapotranspiration (ET)-unit map of Basin and Range carbonate-rock aquifer system study area, Nevada and Utah.

[NA, not applicable]

ET-unit name	Narrative	Accuracy, in percent correct	
		Producer	User
Open water	The number of reference sites is small. The unit generally mapped well.	<sup>1</sup> 75/NA	<sup>1</sup> 100/NA
Marshland	The number of reference sites is small in Nevada. In Utah, the number of reference sites is significantly larger. Confusion occurs with riparian woodland and shrubland.	<sup>1</sup> 18/93	<sup>1</sup> 25/100
Meadowland	Meadowland generally mapped well. The primary source of confusion was with grassland.	<sup>2</sup> 88	<sup>2</sup> 88
Grassland	The number of reference sites was small. The greatest confusion was with meadowland and sparse desert shrubland. Most of the conflicts correlated with the presence of a small annual forb known as <i>halogeton glomeratus</i> . Its presence was likely caused by above normal precipitation during the year.	<sup>2</sup> 33	<sup>2</sup> 33
Moist bare soil	Only Yelland Dry Lake was visited. At the time of the visit, conditions were extremely dry. The playa is known to be persistently moist and the depth to ground water ranges from about 4 to 10 feet (Todd Mihevc, Desert Research Institute, oral commun., 2006).	NA	NA
Dense desert shrubland	The number of reference sites was large. Comparison of mapped and observed conditions indicates that the unit mapped well.	<sup>2</sup> 100	<sup>2</sup> 53
Moderately dense desert shrubland	The number of reference sites was large. In general, the unit mapped well. Limited confusion occurs with sparse desert shrubland.	<sup>2</sup> 69	<sup>2</sup> 72
Sparse desert shrubland	The number of reference sites was adequate. Confusion occurs with moderately dense desert shrubland.	<sup>2</sup> 67	<sup>2</sup> 94
Dry playa	The number of reference sites in Nevada was small. In Utah, the number of reference sites was large. The ET unit was mapped well in Utah. Confusion occurs with salt desert scrub and greasewood flats.	<sup>1</sup> 25/73	<sup>1</sup> 33/25
Xerophyte	The number of reference sites was small.	<sup>2</sup> 50	<sup>2</sup> 50
Recently irrigated cropland	All irrigated fields (cropland) were checked and verified in the field. Misidentifications were corrected accordingly and assumed to be 100 percent accurate.	<sup>3</sup> 100	<sup>3</sup> 100

<sup>1</sup>Source is from SWReGAP, Nevada and Utah. First percent is for Nevada and second percent is for Utah.

<sup>2</sup>Source is from USGS and DRI field visits in Spring and Snake Valleys.

<sup>3</sup>Source is from USGS field visits in all valleys.

The overall accuracy of the ET units delineated from the MSAVI and Tasseled Cap procedures was 72 percent. In general, a number of errors occurred between ET units of one density with an ET unit of the next higher or lower density. For example, an area in the field of sparse desert shrubland was classified as moderately dense desert shrubland. This error can occur because of the difficulty to measure the percentage of vegetation cover. Another example of error is an area that in the field was observed to be sparse desert shrubland was mapped as the ET unit grassland. This probably was the result of an annual plant that was alive during the satellite overpass in the summer, but dormant during the field sampling. A more-accurate map describing the spatial distribution of desert shrub and grass ET units could be developed by using Landsat imagery acquired at different dates within a year and imagery acquired during different years.

## Summary

The purpose of this report is to document the ET-unit map used to delineate areas of vegetation, open water, and moist soil that contribute to ET in the study area. This data set is referred as an ET-unit map.

The data sets used to develop the ET-unit map are based on Landsat satellite data. The SWReGAP program developed a five-state land cover map base on ecological systems. The ecological systems describing open water, marshlands, and dry playa were used to delineate those ET units. Landsat data were acquired for the dates of July 3 and 12 of 2005. To delineate ET units of shrubs and grasses, the Landsat data were used to produce the MSAVI, an index that relates to the greenness and density of plant growth. The ET units of meadowland, grassland, dense desert shrubland, moderately dense desert shrubland, and sparse desert shrubland were delineated. The 2005 imagery also was used to produce the Tasseled Cap transformation. The brightness, greenness, and moisture bands were used to identify ET units of moist bare soil. The 2005 imagery, along with Landsat imagery from 2002 and 2000, was used to identify recently irrigated croplands by photo interpretation.

The final ET-unit map was assembled from the SWReGAP, MSAVI, Tasseled Cap datasets, the interpretations from Landsat imagery and field mapping. The recently irrigated cropland was used as is and had the highest priority. The SWReGAP classes also were used as is and had the second highest priority. The moist bare soil derived from the Tasseled Cap transformation was used and had the next priority. Finally, the ET units from the MSAVI were used at the lowest priority. The MSAVI and Tasseled Cap data sets were filtered such that the minimum mapping unit was about 0.8 acre. The SWReGAP data had a minimum mapping unit of 1.1 acre. The boundary of potential areas of ground-water discharge was used as a mask allowing only those areas within the boundary to be used in the final ET-unit map.

Phreatophytic, xerophytic, and recently irrigated cropland in the study area identified from the ET-unit map have been summarized. About 1,675 mi<sup>2</sup> in the study area are covered with phreatophytes or moist bare soil that potentially discharged ground water. This area does not include acreage of recently irrigated croplands. An accuracy assessment of the ET units was conducted by field visits to Snake and Spring Valleys and has been summarized. The accuracy of the shrubland ET units, accounting for about 83 percent of the potential ground-water discharge acreage, indicated that 67 to 100 percent of the field sites visited was mapped correctly. The ET-unit map developed used a consistent approach to delineate ET units throughout the study area. The ET-unit map also included selected classes from a national land cover data set.

## Acknowledgments

The authors would like to express their appreciation to the Bureau of Land Management and many private landowners in the area that willingly provided access to their property. The authors also acknowledge the effort of the Desert Research Institute in assisting in remote sensing and field-sampling efforts and the Southern Nevada Water Authority for their cooperation in many discussions and the sharing of information pertinent to this effort. The authors would also like to extend their appreciation to William D. Nichols (recently deceased) of the U.S. Geological Survey who recognized the potential of remote sensing in quantifying evapotranspiration in arid environments of the southwestern United States.